

Adelic methods in algebraic geometry

April 11, 2016

1 Motivation

In order to motivate the concepts we will encounter in these notes, and simultaneously test their applicability, we will take a closer look at the interplay of Tate vector spaces and the theory of algebraic curves. We will see how the formalism of Tate objects and adèles can be used to shed light on various classical results and constructions. The Riemann-Roch theorem, and a zoo of classical reciprocity laws (such as the sum of residues theorem) will follow from entirely elementary considerations about infinite-dimensional vector spaces.

This section closely follows Dustin Clausen's treatment [Cla], which was also the authors' port of entry to this fascinating subject. We strongly recommend Clausen's text, particularly to the reader wishing to see applications of this philosophy to the geometric Langlands programme.

1.1 Central extensions of loop groups

Consider the compact Lie group $G = U(n)$. The associated loop group LG is defined to be infinite-dimensional Lie group of smooth maps $S^1 \rightarrow G$. Its Lie algebra $L\mathfrak{g}$ is the vector space of smooth maps $S^1 \rightarrow \mathfrak{g}$. A central construction in the theory of loop groups, as presented in [PS86], is its canonical central extension

$$1 \rightarrow S^1 \rightarrow \widehat{LG} \rightarrow LG \rightarrow 1.$$

Central extensions $1 \rightarrow A \rightarrow \widehat{\Gamma} \rightarrow \Gamma \rightarrow 1$ can be obtained as follows: at first one constructs an action of Γ on X , which carries a transitive and faithful action of BA (in other words is a BA -torsor). We can either think of BA as a homotopy type, that is, really the classifying space of A , or, if A is a discrete group, realise it as the groupoid with a single object which has A as group of automorphisms.

If the action of Γ on X intertwines with the BA -action giving the torsor structures, we have actually constructed a homomorphism $\Gamma \rightarrow BA$, which classifies a central extension of Γ by A .

In the case of the loop group LG , one notices that there is a canonical central extension of the Lie algebra

$$0 \rightarrow \mathbb{R} \rightarrow \widehat{L\mathfrak{g}} \xrightarrow{p} L\mathfrak{g} \rightarrow 0,$$

which corresponds to an alternating pairing $b(-, -): L\mathfrak{g} \times L\mathfrak{g} \rightarrow \mathbb{R}$, such that we have $[x, y]_{\widehat{L\mathfrak{g}}} = [p(x), p(y)] + b(x, y) \cdot \mathbf{1}$. In the case of a loop group one then chooses b to be (up to a constant) equal to

$$\int_{S^1} \langle f, dg \rangle = \int_{S^1} \langle f, g' \rangle d\theta.$$

The definition already indicates a close connection to geometric concepts such as monodromy and the residue pairing. The latter is one of the reasons why it seems feasible to study analogous constructions purely within the realm of algebraic geometry.

Lemma 1.1. *Let M be a simply-connected, connected manifold. Given a closed 2-form ω , such that $\frac{\omega}{2\pi}$ represents an integral cohomology class, there exists a unitary line bundle with a connection (L, ∇) , such that $F(\nabla) = \omega$. Moreover for two such line bundles (L_1, ∇_1) and (L_2, ∇_2) , there exists an isomorphism, unique up to multiplication an element of $U(1)$.*

One applies this lemma to X being the identity component of the homogeneous space LG/T . It is clear that the identity component of LG acts on X , and therefore also on the space of unitary line bundles with connections equal to ω . Hence we are in the situation discussed above, and we see at least how a canonical central extension of the identity component is constructed.


1.2 Tate vector spaces

For the rest of this section we fix a field k , which we assume to be algebraically closed to simplify the exposition. We give a preliminary definition of Tate vector spaces, which we will revisit later in a more general context.

Definition 1.2. A Tate vector space V is a pair (V, Gr_V) , consisting of a k -vector space V , and a set of subsets $\text{Gr}_V = \{L \subset V\}$, satisfying the conditions (A1-4) below. Subsets belonging to Gr_V will be called lattices.

(A1) We have $\bigcap_{L \in \text{Gr}_V} L = \{0\}$, and $\bigcup_{L \in \text{Gr}_V} L = \{V\}$.

(A2) For $L_1, L_2 \in \text{Gr}_V$ there exist lattices $M, N \in \text{Gr}_V$, such that



with the respective subquotients being finite-dimensional

(A3) If $L_1 \subset M \subset L_2$, such that $L_1, L_2 \in \text{Gr}_V$, then $M \in \text{Gr}_V$.

(A4) The natural map

$$V \longrightarrow \varprojlim_{L \in \text{Gr}_V} V/L$$

is an isomorphism.

A familiar example of such a structure is given by the vector space $F = k((t))$ of formal Laurent series. Consider the collection Gr'_F of subspaces $t^n k[[t]]$ for $n \in \mathbb{Z}$. It is clear that the axioms (A1,2,4) are satisfied, so we simply define Gr_V as the saturation or sandwich-closure of Gr'_F .

The fourth axiom indicates that Tate vector spaces shouldn't be viewed as plain vector spaces, but rather as topological vector spaces. Endowing each of the quotients V/L with the discrete topology, and V with the resulting inverse limit topology, we obtain Lefschetz's *linearly compact vector spaces*. We call this functor from the category of Tate spaces to topological vector spaces the *topological realisation*, and denote it by V^{top} .

Definition 1.3. A topological vector space is called *linearly locally compact*, if it can be obtained as the topological realisation of a Tate space.

This construction is as good as it gets. Topological realisation is fully faithful, that is we can reconstruct the pair (V, Gr_V) from the associated linearly locally compact vector space. Speaking from an aesthetic perspective it is relieving that the existence of a collection of lattices Gr_V is a mere property of a topological vector space, rather than an extra structure imposed on V . Let's see why this is true, and how topology allows one to recover the image of lattices L in the topological realisation V^{top} .

Definition 1.4. A topological vector space is called *linearly locally compact*, if it is equivalent to the inverse limit of finite-dimensional (discrete) vector spaces.

Linearly compact spaces W are precisely the topological realisations of Tate spaces V , where $V \in \text{Gr}_V$ is itself a lattice. Indeed, we just let V be the underlying vector space of W , and stipulate $L \subset V$ to belong to Gr_V , if it arises as the kernel of a continuous surjection $V \twoheadrightarrow V/L$ with V/L a finite-dimensional discrete vector space.

If $W = V^{\text{top}}$ is a linearly locally compact vector space, we can reconstruct (V, Gr_V) by forgetting the topology on W , and stipulating that a subspace $L \subset V$ is a lattice, if and only if the associated subspace of V is linearly compact, and agrees with the kernel of a surjection onto a discrete vector space V/L .

Definition 1.5. A Tate vector space V is called discrete, if $\{0\} \in \text{Gr}_V$ is a lattice. This happens precisely when V^{top} is discrete in the topological sense.

Let's imagine for a second that our base field k is finite. In this case, linearly compact vector spaces are actually compact as topological spaces, because they are obtained as inverse limit of finite sets, and similar Tate spaces have locally compact topological realisations. A subspace of a linearly locally compact is a lattice if and only if it is compact and co-discrete (that is, V/L is discrete).

The next lemma claims that every continuous homomorphism from a linearly compact space to a discrete one, factors through a finite-dimensional subspace. In the case of a finite field this is clear, because continuous images of compact spaces are compact, and compact subsets of discrete spaces are finite.

Lemma 1.6. Every morphism $L \rightarrow D$ from a linearly compact Tate space L to a discrete Tate space D factors through a finite-dimensional subspace of D .

As an immediate consequence we obtain that lattices are commensurable in the sense that they only differ by finite-dimensional corrections (which is also a consequence of (A2) of our definition of Tate vector spaces).

Remark 1.7. Let $M \subset N$ be an ordered pair of lattices in a Tate space, then the quotient L_2/L_1 is finite-dimensional.

We define the relative dimension of two lattices $L_1, L_2 \in \text{Gr}_V$ to be the difference

$$\dim(L_1 - L_2) = \dim(L_1/N) - \dim(L_2/N) = \dim(M/L_2) - \dim(M/L_1),$$

where M, N are common upper, respectively lower bounds, which are guaranteed to exist by axiom (A2).

1.3 Torsors and central extensions

Every Tate vector space V gives rise to a whole zoo of torsors, acted on by $\text{Aut}(V)$, the group of continuous automorphisms of V . As a simple warm-up exercise we define the \mathbb{Z} -torsor of dimension theories.

Definition 1.8. A dimension theory is a map $d: \text{Gr}_V \rightarrow \mathbb{Z}$, assigning an integer to every lattice, such that for every pair of lattices we have $d(L_1) - d(L_2) = d(L_1 - L_2)$. The set of dimension theories for V will be denoted by $\text{Dim}(V)$.

Since any two lattices in V are commensurable, we see that for every lattice L , and integer $n \in \mathbb{Z}$ there exists precisely one dimension theory d , satisfying $d(L) = n$. This implies that $\text{Dim}(V)$ is a non-empty set, and that the natural \mathbb{Z} -action $d + k: L \mapsto d(L) + k$ endows $\text{Dim}(V)$ with the structure of a \mathbb{Z} -torsor. Since $\text{Aut}(V)$ acts on the set Gr_V of lattices, we obtain an induced action on $\text{Dim}(V)$.

A dimension theory should be viewed as a *renormalisation* of the infinite-dimensional object V . In general it is not possible to speak of the dimension of a lattice $L \subset V$ in the classical sense. Instead of worrying about definition, we choose the dimension of a lattice arbitrarily, and only ask for internal consistency of this assignment. Such renormalisations play a role in Kontsevich's theory of motivic integration, and in the work of Kapranov–Vasserot [KV07] about categories of D -modules on infinite-dimensional varieties.

As we observed in the first subsection, this gives rise to a map $\nu: \text{Aut}(V) \rightarrow \mathbb{Z}$. We can compute the value of ν on an automorphism f as the relative dimension $\dim(gL - L)$, for an arbitrary lattice $L \in \text{Gr}_V$. The group of units of the field $F = k((t))$ acts on the Tate space F by automorphisms, and therefore we obtain an induced map $\nu: F^\times \rightarrow \mathbb{Z}$. This is the valuation map, which is characterised by the property that $t^{-\nu(f)}f \in k[[t]]^\times$.

The choice of a dimension theory for a Tate space is often compared to the choice of a Haar measure for a local field F . Just like for dimension theories, there is no canonical choice of a Haar measure on F . And if we pullback a measure μ via the multiplication map $x \mapsto ax$, we obtain $a^*\mu = q^{\nu(a)}\mu$, where q denotes the cardinality of the residue field of F .

Let's take this to the next level. This time we put the emphasis on top exterior powers of vector spaces, rather than their dimension. This is sensible because determinants behave formally similar to dimensions. For every short exact sequence $0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$ of finite-dimensional vector spaces we have a natural isomorphism

$$\det(V) \simeq \det(U) \otimes \det(W), \tag{1}$$

in particular we have $\det(U \oplus W) \simeq \det(U) \otimes \det(W)$.

Definition 1.9. A determinant theory assigns to every lattice $L \in \text{Gr}_V$ a line (that is, a 1-dimensional vector space) $\mathcal{L}(L)$, such that for every ordered pair of lattices $M \subset N$, we have $\mathcal{L}(N) \simeq \mathcal{L}(M) \otimes \det(N/M)$, which satisfies for every ordered triple $L \subset M \subset N$ a compatibility condition

$$\begin{array}{ccc} \mathcal{L}(N) & \longrightarrow & \mathcal{L}(M) \otimes \det(N/M) \\ \downarrow & & \downarrow \\ \mathcal{L}(L) \otimes \det(N/L) & \longrightarrow & \mathcal{L}(L) \otimes \det(N/M) \otimes \det(M/L). \end{array}$$

We denote the category (or rather groupoid) of determinant theories by $\text{Det}(V)$. The same argument as in the case of dimension theories shows that a determinant theory is uniquely (up to a unique isomorphism) characterised by the line associated to a single lattice $L \in \text{Gr}_V$. In particular, we see that $\text{Det}(V)$ is a torsor over the groupoid of lines Bk^\times . The natural action of $\text{Aut}(V)$ on Gr_V induces therefore as before an action on $\text{Det}(V)$, which gives rise to a homomorphism $\text{Aut}(V) \rightarrow k^\times$. The corresponding central extension of $\text{Aut}(V)$,

$$1 \rightarrow k^\times \rightarrow \widehat{\text{Aut}(V)} \rightarrow \text{Aut}(V) \rightarrow 1$$

is the promised algebraic analogue of the canonical central extension of loop groups. Indeed, we can embed $L\text{GL}_n = \text{GL}_n((t))$ in the automorphism group of the Tate vector space F^n , and thus obtain a central extension of $L\text{GL}_n$ by pullback.

Just like the dimension torsor $\text{Dim}(V)$ gives rise to a numerical invariant $\nu(f)$ of an automorphism $f \in \text{Aut}(V)$, the central extension induced by $\text{Det}(V)$ can be used to attach an element of k^\times to a pair of commuting automorphisms of V .

Lemma 1.10. Let $1 \rightarrow A \rightarrow \widehat{\Gamma} \xrightarrow{p} \Gamma \rightarrow 1$ be a central extension, and $f, g \in \Gamma$ to commuting elements. Then, $[p^{-1}(f), p^{-1}(g)] \in A$ is a well-defined element of A . If Γ is commutative, we denote the pairing $\Gamma \times \Gamma \rightarrow A$ by $(-, -)_{\widehat{\Gamma}}$.

In the case of the central extension of $L\mathbb{G}_m$, the resulting pairing is a well-known construction in number theory.

Lemma 1.11. The pairing $(-1)^{\nu(f)\nu(g)}(f, g)_{L\mathbb{G}_m}^{-1}$ agrees with the tame symbol

$$(-1)^{\nu(f)\nu(g)} \frac{f^{\nu(g)}}{g^{\nu(f)}} \Big|_{t=0} \in k^\times.$$

The modification of the commutator pairing by the sign $(-1)^{\nu(f)\nu(g)}$ can be explained in a more natural way, by replacing determinants by graded determinants.

Definition 1.12. A graded line is a pair (L, n) , where L is a 1-dimensional space, and $n \in \mathbb{Z}$ is an integer. The graded determinant of a finite-dimensional vector space V is the pair $(\det V, \text{rk } V)$.

The tensor product of two graded lines $(M, m), (N, n)$ is simply defined by the tensor product of the two lines, and addition of integers. However, we modify the natural isomorphism $M \otimes N \xrightarrow{\simeq} N \otimes M$ by multiplication with the sign $(-1)^{mn}$.

By consistently adding the word *graded* to Definition 1.9 one obtains the definition of a *graded determinant theory*.

1.4 Adèles

The time has come to see the theory of Tate objects in action in a global situation. Let X be a smooth curve over $k = \bar{k}$ (sometimes, we will assume X to be projective). For every line E/X we define the vector space of adèles to be

$$\mathbb{A}_X(E) = \prod'_{x \in X_{cl}} (E_x \otimes \widehat{F}_x) = \{(f_x)_{x \in X_{cl}} \mid f_x \in E_x \otimes \widehat{\mathcal{O}}_x \text{ for almost all } x\}.$$

For $E = \mathcal{O}_X$ the trivial line bundle, we obtain a ring, denoted by \mathbb{A} . In general, $\mathbb{A}(E)$ is an \mathbb{A} -module.

However, $\mathbb{A}_X(E)$ also carries the structure of a Tate vector space. The collection of lattices is given by the saturation of $\text{Gr}'_{\mathbb{A}(E)}$, consisting of the subspaces

$$\mathbb{O}_D(E) = \{(f_x)_{x \in X_{cl}} \mid f_x \in t^{-\nu_x(D)}(E_x \otimes \widehat{\mathcal{O}}_x) \text{ for almost all } x\},$$

indexed by divisors on X . We denote $\mathbb{O}_0(E)$ also by $\mathbb{O}(E)$, and refer to it as integral adèles.

Theorem 1.13 (Weil). *There is a canonical equivalence $\text{Pic}(X) \cong [\mathbb{G}_m(F) \setminus \mathbb{G}_m(\mathbb{A}) / \mathbb{G}_m(\mathbb{O})] = [F^\times \setminus \mathbb{A}^\times / \mathbb{O}^\times]$.*

Weil's theorem holds in fact for a wide range of groups. For example we can describe $\text{Bun}_n(X)$, the stack of rank n bundles on X , as the double quotient $[\text{GL}_n(F) \setminus \text{GL}_n(\mathbb{A}) / \text{GL}_n(\mathbb{O})]$. The case of line bundles however, is easy to prove, since $\mathbb{A}^\times / \mathbb{O}^\times$ is equivalent to the free abelian group generated by the closed points of X , that is, the group of divisors Div_X . The corresponding action of F^\times on Div_X takes a divisor D and modifies it by the principal divisor associated to a non-zero invertible function f . Therefore, we see that Weil's theorem for line bundles is just another way to state the classical identity

$$\text{Pic}(X) \cong \text{Div}_X / \text{Div}_X^{\text{princ}},$$

that line bundles on X can be presented by divisors modulo principal divisors.

Lemma 1.14. *Let $f \in \mathbb{A}^\times$ be an invertible adèle (also known as an idèle). The order $\nu(f)$ computed with respect to $\text{Dim}(\mathbb{A})$ agrees with the degree $\deg(f) = \deg(D)$ of the divisor $D \in \text{Div}_X = \mathbb{A}^\times / \mathbb{O}^\times$ associated to f .*

Indeed, we just have to choose a dimension theory d , and compute the difference $d \circ f - d$. If we choose d to be the unique dimension theory which assigns to the lattice $\mathbb{O} \subset \mathbb{A}$ the integer 0, we see directly that $\nu(f)$ is equal to the relative dimension $\dim(f\mathbb{O} - \mathbb{O}) = \deg(f)$.

We could have chosen a completely different dimension theory, and still have obtained the same integer $\nu(f)$. Another interesting choice for a projective curve X is

$$\chi: \mathbb{O}_{D'} \mapsto \chi(\mathcal{O}(D')) = \dim H^0(X, \mathcal{O}(D')) - \dim H^1(X, \mathcal{O}(D')).$$

The long exact sequence in cohomology implies readily that χ is a dimension theory, and we see that $\nu(f) = \chi(\mathcal{O}(D)) - \chi(\mathcal{O})$, which can be summarised as the assertion of the Riemann-Roch theorem

$$\chi(\mathcal{O}(D)) - \chi(\mathcal{O}) = \deg D.$$

Replacing the dimension torsor by $\text{Det}(V)$, or rather its graded analogue, one obtains the following multiplicative analogue of the Riemann-Roch theorem:

Theorem 1.15 (Weil reciprocity). *Let f, g be two non-zero rational functions on a projective curve X , then we have that the product below is well-defined, and $\prod_{x \in X_{cl}} (-1)^{\nu(f)\nu(g)} \frac{f^{\nu(g)}}{g^{\nu(f)}}(x) = 1$.*

Let us explain how this works. We denote the field of rational functions on X by K . The simple observation that two rationally equivalent divisors have the same cohomology groups, implies that the central extension of K^\times , obtained from the action on \mathbb{A}^\times is trivial. Hence, the commutator pairing is equal to 1, for elements of K^\times .

1.5 An adelic viewpoint on cohomology

When discussing the proofs of Riemann-Roch and Weil reciprocity, using adèles, we were making use of cohomology theory, which on a priori seems to be external to the theory of adèles. However, one can use adèles to compute cohomology groups.

Proposition 1.16. *Let E be a coherent sheaf on X , and $\mathbb{A}(E)$, $\mathbb{O}(E)$, and $K(E) = K \otimes_{\mathcal{O}} E$ be the abelian groups of adèlic, integral adèlic, and rational sections. Then, we have that the cohomology groups of the complex*

$$\mathbb{O}(E) \oplus F(E) \xrightarrow{\text{incl} - \text{incl}} \mathbb{A}(E)$$

are canonically equivalent to $H^i(X, E)$. In fact, this complex is quasi-isomorphic to the complex of derived global sections $R\Gamma(X, E)$.

This proposition is easy to prove. At first one observes that replacing X by a Zariski open subset, one obtains a complex of flasque sheaves, which happens to be a resolution of E . The latter property is a consequence of the tautology that a rational section of E , without poles, is a regular section.

The complex above is quasi-isomorphic to

$$[\mathbb{O}(E) \longrightarrow \mathbb{A}(E)/F(E)], \tag{2}$$

and using this presentation one can give an adelic proof of Serre duality.

Theorem 1.17 (Adelic Serre duality). *For a smooth projective curve, the residue pairing $\text{res}: \mathbb{A} \times \mathbb{A}(\Omega_X^1) \longrightarrow \mathbb{A}$ is perfect, that is induces an equivalence between $\mathbb{A}(\omega_X)$ and the (topological) dual of \mathbb{A}_X . In particular we see that $\mathbb{A}(E)^\vee \cong \mathbb{A}(E^\vee \otimes \Omega_X^1)$. These isomorphisms descend to isomorphisms $(\mathbb{A}(E)/\mathbb{O}(E))^\vee \cong \mathbb{O}(E)^\vee$, and $F(E)^\vee \cong \mathbb{A}(E^\vee \otimes \Omega_X^1)/F(E^\vee \otimes \Omega_X^1)$.*

The first two assertions can be checked by a direct computation, and actually don't rely on projectivity. The third assertion does use projectivity. Even just to define a pairing, we use the residue formula for rational 1-forms.

Using the adelic duality theorem above, we obtain a proof of Serre duality, because the (topological) dual of the complex (2) is now equivalent to

$$[F(E^\vee \otimes \Omega_X^1) \longrightarrow \mathbb{A}(E^\vee \otimes \Omega_X^1)/\mathbb{O}(E^\vee \otimes \Omega_X^1)],$$

which is just another way to write down the complex of Proposition 1.16 for $E^\vee \otimes \Omega_X^1$.

2 From adèles for curves to multidimensional adèles

In this section we will discuss the transition from the 1-dimensional case to higher dimensions. This is inspired by higher class field theory (established by Kato and Saito).

2.1 Classical story: function field case

We fix X/k a classical curve, and E a line bundle on X . The integral adèles are defined to be

$$\mathbb{O}(E) = \prod_{x \in X_0} \widehat{\mathcal{O}}_x \otimes E,$$

and the adèles are defined to be

$$\mathbb{A}_E = \text{colim}_D \mathbb{O}_{\mathcal{O}_D \otimes (E)} = \prod'_{x \in X_{cl}} (E_x \otimes \widehat{F}_x) = \{(f_x)_{x \in X_{cl}} \mid f_x \in E_x \otimes \widehat{\mathcal{O}}_x \text{ for almost all } x\}.$$

Over a finite fields, these are compact abelian topological groups, respectively locally compact abelian topological groups, simply because $k[[t]]$ and $k((t))$ has this property. The modern viewpoint is to work functorially, that is replace every appearance of factors such as $k((t))$ by the functor, sending a commutative ring R to $R((t))$. An equivalent perspective is to work with Tate vector spaces, respectively bundles.

Recall the archetypical example of a Tate vector space: the space of formal Laurent series $F = k((t))$ over a field k . We defined Gr_F as the saturation $\{t^n k[[t]] \mid n \in \mathbb{Z}\}'$. Similarly one defines $\text{Gr}_{\mathbb{A}_E} = \{\mathbb{O}_{\mathcal{O}_D} \otimes E\}'$, which endows the adèles with the structure of a Tate space.

Recall that we discussed the notion of a dimension theory, which was a map $\text{Gr}_V \rightarrow \mathbb{Z}$, consistent with relative dimension of lattices.

In the case of F , we obtained a dimension torsor $\text{Dim}(F)$, such that the resulting map $k((t))^\times \xrightarrow{\nu} \mathbb{Z}$ agrees with the t -valuation of F .

Let's discuss the analogous situation for the adèles: we saw that $\text{Pic}(X) \cong [k(X)^\times \setminus \mathbb{A}^\times / \mathbb{O}^\times]$, and the dimension torsor of the Tate object \mathbb{A} induced a map $\text{deg}: \text{Pic}(X) \rightarrow \mathbb{Z}$, which agrees with the degree of line bundles. We then used this observation to reprove the Riemann-Roch formula for curves. The key step to make this connection was to use the dimension theory given by the Euler characteristic of the cohomology of line bundles. Now that we've seen the applicability of adèles we'll try to generalise this theory to higher-dimensions.

2.2 Adèles for schemes (after Parshin, Beilinson)

In the case of curves, adèles were realised as a sort of restricted product, ranging over closed points of X , or the places of $k(X)$. In the higher-dimensional case, their place will be taken by flags of irreducible subsets.

The local factors for a surface, are expected to be of the shape $\widehat{\mathcal{O}_{\eta_2, \eta_1}}$, where η_1 is the generic point of a curve, containing a closed point η_2 (the subscript encodes the codimension of the closure of a point).

The local factors have the structure of an n -local field.

$$\begin{array}{ccc}
 k((s))((t)) & & \\
 \uparrow & & \\
 k((s))[[t]] & \twoheadrightarrow & k((s)) \\
 & & \uparrow \\
 & & k[[t]] \twoheadrightarrow k
 \end{array}$$

The next step is to take some case of restricted product. However, there's a recursive definition, which helps us avoiding a lot of guesswork what the right definition should be.

Let's what kind of structure we can put on higher-dimensional adèles. We'll work with the local factors, in order to get a feeling for this problem. Consider a 2-local field, such as $K = k((s))((t))$. Is there a well-behaved topology on it?

Example 2.1. *Let \mathcal{T} be any topology on K , such that $k((s))[[t]] \twoheadrightarrow k((s))$ is a quotient topology map, then \mathcal{O}_K (the ring of integers), is not a topological ring.*

This is disappointing. Yekutieli has therefore studied adèles using the theory of semi-topological rings. Fesenko proposed to work with sequential topological rings, however these approaches are not directly compatible. Kato suggested that topology is simply not the right structure to study higher local fields, and proposed to work with categories of iterated Ind-Pro objects. This works, because $k((s))$ can be defined as

$$\text{colim}_j \lim_i s^{-j} k[s]/(s^i).$$

Remark 2.2. *Pro-systems carry more information than their topological realisations. There are pro-systems of sets, such that the resulting space is empty, but the pro-system carries a lot of information which would be lost if we didn't retain it in a formal manner.*

2.3 Dimension theories for 2-local fields

We fix an equicharacteristic 2-local field $K = k((s))((t))$, and choose a splitting $k((s))[[t]] \oplus t^{-1}k((s))[t^{-1}]$. The problem is that a general automorphism won't respect such a splitting.

Subquotients of lattices (with respect to the variable t), will look like $t^2/k((s))[[t]]/t^{10}k((s))[[t]] \cong k((s))^{\oplus 8}$, which is infinite-dimensional over the base field k . In order to rectify this, we also have to consider s -lattices.

Picture

add picture

This shows that for 2-local fields the natural construction is to work with pairs of automorphisms. In the definition below we denote by \mathcal{O}_1 the ring $k((s))[[t]]$. The definition below is intentionally incomplete, and will be corrected below.

Definition 2.3 (Beilinson, Yekutieli). *Suppose V is a finite-dimensional K -vector space, a lattice is a finitely generated \mathcal{O}_1 -submodule of V , such that $K \cdot L = V$. We define $E^{Yek}(V_1, V_2)$ (the analogue of continuous homomorphisms) to be the vector space of k -linear maps $\phi: V_1 \rightarrow V_2$, such that for every pair of lattices $L_1 \subset V_1$, $L_2 \subset V_2$, there exist lattices $L'_1 \subset V_1$ and $L'_2 \subset V_2$, such that $L'_1 \subset L_1$, $L_2 \subset L'_2$, and $\phi(L'_1) \subset L_2$, and $\phi(L_1) \subset L'_2$, and with the resulting maps $\bar{\phi}: L_1/L'_1 \rightarrow L'_2/L_2$ belonging to $E^{Yek}(L_1/L'_1, L'_2/L_2)$.*

The problem with this definition is that in order for the last condition to make sense, we have to be able to consider the subquotients L_1/L'_1 , and L'_2/L_2 as modules over $k((s))$. This is possible once we've chosen splittings.

3 Beilinson-Parshin adèles

Adèles were first introduced by Parshin for surfaces in 1976, and for Noetherian schemes by Beilinson in 1980. Let X be a Noetherian scheme, and $\eta_0, \eta_1 \in X$. We define $\eta_0 \geq \eta_1$ to be equivalent to $\eta_1 \in \overline{\{\eta_0\}}$.

Suppose $K \subset S(X)_\eta$, define ${}_\eta K = \{(\eta_1 \geq \dots \geq \eta_m) | \eta \geq \eta_1 \geq \dots \geq \eta_m \in K\}$.

If \mathcal{F} is a coherent sheaf on X , define for

- $n = 0$: $A(K, \mathcal{F}) = \prod_{\eta \in K} \varprojlim (\mathcal{F} \otimes \mathcal{O}_\eta / \mathfrak{m}_\eta^i)$.
- $n \geq 1$: $A(K, \mathcal{F}) = \prod_{\eta \in X} \varprojlim_i A({}_\eta K, \mathcal{F} \otimes \mathcal{O}_{X, \eta} / \mathfrak{m}_\eta^i)$,

and if \mathcal{F} is quasi-coherent, we stipulate $A(K, \mathcal{F}) = \text{colim}_i A(K, \mathcal{F}_i)$ with $\text{colim } \mathcal{F}_i \cong \mathcal{F}$ a colimit of coherent sheaves. By induction this defines a unique family of functors $A(K, -)$ from quasi-coherent sheaves on X to \mathcal{O}_X -modules. We are mostly interested in the values of these functors for coherent sheaves, however note that we cannot unravel the induction without including the case of quasi-coherent sheaves in the definition above.

Example 3.1. *We fix a curve $X = \text{Spec } k[s, t]/(s^3 + s^2 - t^2)$, which we allow to be singular. Let Δ be the flag $(0) \geq (s, t)$. Since we have $s^3 + s^2 - t^2 = (s\sqrt{1+s} + t)(s\sqrt{1+s} - t)$, one sees that $A(\Delta, \mathcal{O}_X) = k((t_1)) \oplus k((t_2))$.*

This is a special case of the following theorem.

Theorem 3.2 (Beilinson, first published proof by Yekutieli). *Let X be a Noetherian reduced excellent scheme of dimension n , and $\Delta = (\eta_0 \geq \dots \geq \eta_n)$ a flag with $\text{codim } \{\eta_i\} = i$, then*

$$A({}_{\eta_0} \Delta, \mathcal{O}_X) \subset \prod \mathcal{O}_1 \subset \prod_{\text{finite}} K_i = A(\Delta, \mathcal{O}_X),$$

where K_i are n -local fields.

We want an efficient calculus of lattices. We follow the suggestion of Kato and Beilinson from 1980 that Tate vector spaces can be emulated using categories of ind-pro objects. We fix an exact category \mathcal{C} , e.g. an abelian category \mathcal{C} .

Definition 3.3. *We define $\text{Ind}^a(\mathcal{C})$ to be the full subcategory of $\text{Ind } \mathcal{C}$ of admissible ind objects, that is, objects which allow a presentation $C_1 \hookrightarrow C_2 \hookrightarrow \dots$ with transition maps being admissible monomorphisms.*

Inverting directions of morphisms, and replacing admissible monomorphisms by admissible epimorphisms, one obtains the definition of $\text{Pro}^a(\mathcal{C})$.

Definition 3.4. *An object $X \in \text{Ind}^a \text{Pro}^a(\mathcal{C})$ is called an elementary Tate object, if it has a lattice, that is there exists $L \in \text{Pro}^a(\mathcal{C})$, and an admissible monomorphism $L \hookrightarrow X$, such that $X/L \in \text{Ind}^a(\mathcal{C})$. We denote the resulting exact category by $\text{Tate}^{\text{el}}(\mathcal{C})$. The idempotent completion is denoted by $\text{Tate}(\mathcal{C})$, and its objects are called Tate objects.*

We define categories of n -Tate objects by iteratively applying the Tate construction. Beilinson's adèles can be promoted to take values in exact categories of these iterated Tate objects.

The endomorphism algebra à la Yekutieli of a higher local field, as we defined it in the last section, is equivalent to the endomorphism algebra of n -Tate objects.

4 Generalised dimension and determinant theories

Define the Sato complex $\mathrm{Gr}_n^{\leq}(\mathbf{C}) = \{(L_0 \hookrightarrow \cdots \hookrightarrow L_n \hookrightarrow V)\}$, where the L_i denote lattices in an elementary Tate object V .

Definition 4.1.

$$\begin{array}{ccc} & \mathrm{Gr}_{\bullet}^{\leq}(\mathbf{C}) & \\ & \swarrow \quad \searrow & \\ \mathrm{Tate}^{\mathrm{el}}(\mathbf{C})^{\times} & & S_{\bullet}\mathbf{C}^{\times}, \end{array}$$

where the map to the left forgets lattices, and the map to the right sends the collection of lattices to $(0 \hookrightarrow L_1/L_0 \hookrightarrow \cdots \hookrightarrow L_n/L_0)$. Taking geometric realisation, we obtain the map $\mathrm{Tate}^{\mathrm{el}}(\mathbf{C})^{\times} \rightarrow BK_{\mathbf{C}}$, respectively $\Omega K_{\mathrm{Tate}^{\mathrm{el}}(\mathbf{C})} \rightarrow K_{\mathbf{C}}$, which are called the index map.

The index map generalises the construction of the dimension torsor, as it produces for every elementary Tate object a torsor over the K -theory space $K_{\mathbf{C}}$, together with an action of $\mathrm{Aut}(V)$. For instance, if $\mathbf{C} = \mathrm{Proj} R$, then we have the rank map from $K_{\mathbf{C}}$ to \mathbb{Z} , and by changing the structures group of torsors, we obtain the dimension torsor $\mathrm{Dim}(V)$.

References

- [Cla] D. Clausen, *Infinite-dimensional linear algebra, etc.* 1
- [KV07] Mikhail Kapranov and Éric Vasserot, *Formal loops. II. A local Riemann-Roch theorem for determinantal gerbes*, Ann. Sci. École Norm. Sup. (4) **40** (2007), no. 1, 113–133. MR 2332353 (2008k:14048) 1.3
- [PS86] Andrew Pressley and Graeme Segal, *Loop groups*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1986, Oxford Science Publications. MR 900587 (88i:22049) 1.1